

Class XII: Physics
Chapter : Nuclei

TOP Concepts

1 **Atomic Number.** The number of protons in the nucleus is called the atomic number. It is denoted by Z.

2 **Mass number.** The total number of protons and neutrons present in a nucleus is called the mass number of the element. It is denoted by A.

Number of protons in an atom = Z

Number of electrons in an atom = Z

Number of nucleons in an atom = A

Number of neutrons in an atom = N = A - Z.

3 **Nuclear Mass.** The total mass of the protons and neutrons present in a nucleus is called the nuclear mass.

4 **Nuclide.** A nuclide is a specific nucleus of an atom characterized by its atomic number Z and mass number A. It is represented as ${}_Z X^A$

where X = chemical symbol of the element.

Z = atomic number, and

A = mass number.

5 **Isotopes.** The atoms of an element which have the same atomic number but different mass number are called isotopes. Isotopes have similar chemical properties but different physical properties.

6 **Isobars.** The atoms having the same mass number but different atomic number are called isobars.

7 **Isotones.** The nuclides having the same number of neutrons are called isotones.

8 **Isomers.** These are nuclei with same atomic number and same mass number but in different energy states.

9 **Electron Volt:** It is defined as the energy acquired by an electron when it is accelerated through a potential difference of 1 volt and is denoted by eV.

$$1 \text{ eV} = 1.602 \times 10^{-19} \text{ J}$$

$$1 \text{ MeV} = 10^6 \text{ eV} = 1.602 \times 10^{-13} \text{ J}$$


10 **Atomic Mass Unit:.** It is $\frac{1}{12}$ th of the actual mass of a carbon atom of isotope ${}_6\text{C}^{12}$. It is denoted by amu or just by u.

$$1 \text{ amu} = 1.660565 \times 10^{-27} \text{ kg}$$


The energy equivalence of 1 amu is

$$1 \text{ amu} = 931 \text{ MeV}$$

11 **Discovery of Neutrons.** Neutrons were discovered by Chadwick in 1932. When beryllium nuclei are bombarded by α -particles, highly penetrating radiations are emitted, which consists of neutral particles, each having mass nearly that of a proton. These particles were called neutrons.



A free neutron decays spontaneously, with a half life of about 900 s, into a proton, electron and an antineutrino.



- 12 **Size of the Nucleus.** It is found that a nucleus of mass number A has a radius

$$R = R_0 A^{1/3}$$

where $R_0 = 1.2 \times 10^{-15} \text{ m}$.

This implies that the volume of the nucleus, which is proportional to R^3 is proportional A.

Density of nucleus is constant; independent of A, for all nuclei and density of nuclear matter is approximately $2.3 \times 10^{17} \text{ kg m}^{-3}$ which is very large as compared to ordinary matter, say water which is 10^3 kg m^{-3} .

- 13 **Mass-Energy equivalence:** Einstein proved that it is necessary to treat mass as another form of energy. He gave the mass-energy equivalence relation as

$$E = mc^2$$

where m is the mass and c is the velocity of light in vacuum.

- 14 **Mass Defect.** The difference between the rest mass of a nucleus and the sum of the rest masses of its constituent nucleons is called its mass defect. It is given by

$$\Delta m = [Zm_p + (A - Z)m_n] - m$$

- 15 **Binding Energy.** It may be defined as the energy required to break up a nucleus into its constituent protons and neutrons and to separate them to such a large distance that they may not interact with each other.

It may also be defined as the surplus energy which the nucleus gives up by virtue of their attractions which they become bound together to form a nucleus.

The binding energy of a nucleus ${}_Z X^A$ is given by

$$\text{B.E.} = [Zm_p + (A - Z)m_n - m]c^2$$

- 16 **Binding Energy per Nucleon.** It is average energy required to extract one nucleon from the nucleus. It is obtained by dividing the binding energy of a nucleus by its mass number.

$$\bar{B} = \frac{\text{B.E.}}{A} = \frac{[Zm_p + (A - Z)m_n - m]c^2}{A}$$

- 17 **Nuclear Forces.** These are the strong in attractive forces which hold protons and neutrons together in a tiny nucleus. These are short range forces which operate over very short distance of about 2 – 3 fm of separation between any two nucleons. The nuclear force does not depend on the charge of the nucleon.

- 18 **Nuclear Density.** The density of a nucleus is independent of the size of the nucleus and is given by

$$\rho_v = \frac{\text{Nuclear mass}}{\text{Nuclear volume}}$$

$$= \frac{m_v}{\frac{4}{3}\pi R^2} = 2.9 \times 10^{17} \text{ kg m}^{-3}.$$

19 **Radioactivity.** It is the phenomenon of spontaneous disintegration of the nucleus of an atom with the emission of one or more radiations like α -particles, β -particles or γ -rays. The substances which spontaneously emit penetrating radiation are called radioactive substances.


20 **Radioactivity Displacement Law.** It states that

(i) When a radioactive nucleus emits an α -particle, atomic number decreases by 2 and mass number decreases by 4.


(ii) When a radioactive nucleus emits β -particle, its atomic number increases by 1 but mass number remains same.

(iii) The emission of a γ -particle does not change the mass number or the atomic number of the radioactive nucleus. γ -particle emission by a radioactive nucleus lowers its energy state.


21 **Alpha Decay.** It is the process of emission of an α -particle from a radioactive nucleus. It may be represented as



22 **Beta Decay.** It is the process of emission of an electron from a radioactive nucleus. It may be represented as



23 **Gamma Decay.** It is the process of emission of a γ -ray photon during the radioactive disintegration of a nucleus. It can be represented as



24 **Radioactive Decay Law.** It states that the number of nuclei disintegrated of undecayed radioactive nuclei present at that instant. It may be written as

$$N(t) = N(0)e^{-\lambda t}$$

where $N(0)$ is the number of nuclei at $t = 0$ and λ is disintegration constant.

25 **Decay or disintegration Constant.** It may be defined as the reciprocal or the time interval in which the number of active nuclei in a given radioactive sample reduces to 36.8% $\left(\text{or } \frac{1}{e} \text{ times} \right)$ of its initial value.

26 **Half-life.** The half-life of a radioactive substance is the time in which one-half of its nuclei will disintegrate. It is inversely proportional to the decay constant of the radioactive substance.

$$T_{1/2} = \frac{0.693}{\lambda}$$

27 **Mean Life.** The mean-life of a radioactive sample is defined as the ratio of the combined age of all the atoms and the total number of atoms in the given sample. It is given by

$$\tau = \frac{T_{1/2}}{0.693} = 1.44T_{1/2}$$

28 **Rate of Decay or Activity of a Radioactive Sample.** It is defined as the number of radioactive disintegrations taking place per second in a given sample. It is expressed as

$$R(t) = \left[\frac{dN}{dt} \right] = \lambda N(t) = \lambda N(0)e^{-\lambda t}$$

29 **Curie.** It is the SI unit of decay. One curie is the decay rate of 3.7×10^{10} disintegrations per second.

$$1 \text{ Ci (curie)} = 3.70 \times 10^{10} \text{ disintegrations/s}$$


30 **Rutherford.** One rutherford is the decay rate of 10^6 disintegrations per second.

31 **Natural Radioactivity.** It is the phenomenon of the spontaneous emission of α -, β - or γ -radiations from the nuclei of naturally occurring isotopes.

32 **Artificial or Induced Radioactivity.** It is the phenomenon of inducing radioactivity in certain stable nuclei by bombarding them by suitable high energy sub atomic particles.

33 **Nuclear Reaction.** It is a reaction which involves the change of stable nuclei of one element into the nucleus of another element.

34 **Nuclear Fission.** It is the process in which a heavy nucleus when excited gets split into two smaller nuclei of nearly comparable masses. For example,



35 **Nuclear Reactor.** It is a device in which a nuclear chain reaction is initiated, maintained and controlled.

36 **Nuclear Fusion.** It is the process of fusion of two smaller nuclei into a heavier nucleus with the liberation of large amount of energy.

37 **Critical size and Critical Mass.** The size of the fissionable material for which reproduction factor is unity is called critical size and its mass is called critical mass of the material. The chain reaction in this case remains steady or sustained.

38 **Moderator.** Any substance which is used to slow down fast moving neutrons to thermal energies is called a moderator. The commonly used moderators are water, heavy water (D_2O) and graphite.